

REMARKS

The claims have been amended to eliminate multiple dependency and to place them in better form for U.S. practice. Favorable action on the application is solicited.

Respectfully submitted,

NOVAK DRUCE DeLUCA & QUIGG, LLP

A handwritten signature in cursive script that reads "Herbert B. Keil".

Herbert B. Keil
Reg. No. 18,967

1300 Eye Street, N.W.
Suite 400 East
Washington, D.C. 20005
(202) 659-0100
HBK/sb


Encl.: Listing of Claims

COMPLETE LISTING OF CLAIMS

1. (original) A monocyclopentadienyl complex which contains the structural feature of the formula (Cp) (-Z-A)_mM (I), where the variables have the following meanings:

Cp is a cyclopentadienyl system,

Z is a bridge between A and Cp and is selected from the group consisting of



where

L^{1B}-L^{3B} are each, independently of one another, carbon or silicon,

R^{1B}-R^{6B} are each, independently of one another, hydrogen, C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₆-C₂₀-aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part or SiR^{7B}₃, where the organic radicals R^{1B}-R^{6B} may also be substituted by halogens and two geminal or vicinal radicals R^{1B}-R^{6B} or a radical R^{1B}-R^{6B} and A may also be joined to form a five- or six-membered ring and


R^{7B} are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl or alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two radicals R^{7B} may also be joined to form a five- or six-membered ring,

A is an unsubstituted, substituted or fused, heteroaromatic ring system,

M is a metal selected from the group consisting of titanium in the oxidation state 3, vanadium, chromium, molybdenum and tungsten and

m is 1, 2 or 3.


2. (original) A monocyclopentadienyl complex as claimed in claim 1 of the formula



where the variables have the following meanings:

Cp is a cyclopentadienyl system,

Z is a bridge between A and Cp and is selected from the group consisting of



where

$L^{1B}-L^{3B}$ are each, independently of one another, carbon or silicon,

$R^{1B}-R^{6B}$ are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part or SiR^{7B}_3 , where the organic radicals $R^{1B}-R^{6B}$ may also be substituted by halogens and two geminal or vicinal radicals $R^{1B}-R^{6B}$ may also be joined to form a five- or six-membered ring and

R^{7B} are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl or alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two radicals R^{7B} may also be joined to form a five- or six-membered ring,

A is an unsubstituted, substituted or fused, heteroaromatic ring system,

M is a metal selected from the group consisting of titanium in the oxidation state 3, chromium, molybdenum and tungsten,

m is 1, 2 or 3,

X are each, independently of one another, fluorine, chlorine, bromine, iodine, hydrogen, C_1-C_{10} -alkyl, C_2-C_{10} -alkenyl,


C₆-C₂₀- aryl, alkylaryl having 1-10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part, NR¹R², OR¹, SR¹, SO₃R¹, OC(O)R¹, CN, SCN, β-diketonate, CO, BF₄⁻, PF₆⁻ or a bulky noncoordinating anion,

R¹-R² are each, independently of one another, hydrogen, C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₆-C₂₀-alkenyl, C₆-C₂₀-aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part, SiR³₃, where the organic radicals R¹-R² may also be substituted by halogens and two radicals R¹-R² may also be joined to form a five- or six-membered ring,

R³ are each, independently of one another, hydrogen, C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₆-C₂₀-aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two radicals R³ may also be joined to form a five- or six-membered ring and

k is 1, 2 or 3.

3. (currently amended) A monocyclopentadienyl complex as claimed in claim 1 or 2, wherein the cyclopentadienyl system Cp has the formula (II):




where the variables have the following meanings:

- $E^{1A}-E^{5A}$ are each carbon or not more than one $E^{1A}-E^{5A}$ is phosphorus,
- $R^{1A}-R^{5A}$ are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part, NR^{6A}_2 , $N(SiR^{6A}_3)_2$, OR^{6A} , $OSiR^{6A}_3$, SiR^{6A}_3 , BR^{6A}_2 , where the organic radicals $R^{1A}-R^{5A}$ may also be substituted by halogens and two vicinal radicals $R^{1A}-R^{5A}$ may also be joined to form a five- or six-membered ring, and/or two vicinal radicals $R^{1A}-R^{5A}$ are joined to form a heterocycle which contains at least one atom from the group consisting of N, P, O and S, and where 1, 2 or 3 substituents $R^{1A}-R^{5A}$ is a group-Z-A and
- R^{6A} are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl radical and 6-20 carbon atoms in the aryl radical and two geminal radicals R^{6A} may also be joined to form a five- or six-membered ring.

4. (currently amended) A monocyclopentadienyl complex as claimed in claim 1 any of ~~claims 1 to 3~~, wherein the cyclopentadienyl system Cp together with-Z-A has


the formula (IV):



where the variables have the following meanings:

- $E^{1A}-E^{5A}$ are each carbon or at most one E^{1A} to E^{5A} is phosphorus,
- $R^{1A}-R^{4A}$ are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part, NR^{6A}_2 , $N(SiR^{6A})_3$, OR^{6A} , $OSiR^{A3}$, SiR^{A3} , where the organic radicals $R^{1A}-R^{4A}$ may also be substituted by halogens and two vicinal radicals $R^{1A}-R^{4A}$ may also be joined to form a five- or six-membered ring, and/or two vicinal radicals $R^{1A}-R^{4A}$ may be joined to form a heterocycle containing at least one atom from the group consisting of N, P, O and S,
- R^{6A} are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl, alkylaryl, having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two geminal radicals R^{6A} may also be joined to form a five- or six-membered ring[.],

A is an unsubstituted, substituted or fused, heteroaromatic ring system,
 Z is a bridge between A and Cp and is selected from the group consisting of





where

L^{1B} - L^{3B} are each, independently of one another, carbon or silicon,

R^{1B} - R^{6B} are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part or SiR^{7B}_3 , where the organic radicals R^{1B} - R^{6B} may also be substituted by halogens and two geminal or vicinal radicals R^{1B} - R^{6B} may also be joined to form a five- or six-membered ring and

R^{7B} are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl or alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two radicals R^{7B} may also be joined to form a five- or six-membered ring.

5. (currently amended) A monocyclopentadienyl complex as claimed in claim 1 ~~any of claims 1 to 4~~, wherein A has the formula (IIIA) or (IIIB) :



where the variables have the following meanings:

$E^{1C}-E^{4C}$ are each carbon or nitrogen,

$R^{1C}-R^{4C}$ are each, independently of one another, hydrogen, C_1-C_{20} -alkyl,

C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10

carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl

part or SiR^{5C}_3 , where the organic radicals $R^{1C}-R^{4C}$ may also be

substituted by halogens or nitrogen and further C_1-C_{20} -alkyl,

C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10

carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl

part or SiR^{5C}_3 groups and two vicinal radicals $R^{1C}-R^{4C}$ or R^{1C}

and Z may also be joined to form a five- or six-membered ring and

R^{5C} are each, independently of one another, hydrogen, C_1-C_{20} -alkyl,

C_2-C_{20} -alkenyl, C_6-C_{20} -aryl or alkylaryl having from 1 to 10

carbon atoms in the alkyl part and 6-20 carbon atoms

in the aryl part and two radicals R^{5C} may also be joined to form

a five- or six-membered ring and


p is 0 when $E^{1C}-E^{4C}$ is nitrogen and 1 when $E^{1C}-E^{4C}$ is carbon,
 G^{1C} is nitrogen, phosphorus, sulfur or oxygen,
 $R^{6C}-R^{8C}$ are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part or SiR^{9C}_3 , where the organic radicals $R^{6C}-R^{8C}$ may also be substituted by halogens or nitrogen and C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part or SiR^{9C}_3 groups and two vicinal radicals $R^{6C}-R^{8C}$ or R^{6C} and Z may also be joined to form a 5- or 6-membered ring and

R^{9C} are each, independently of one another, hydrogen C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl or alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two radicals R^{9C} may also be joined to form a five- or six-membered ring and

g is 0 when G^{1C} is sulfur or oxygen and 1 when G^{1C} is nitrogen or phosphorus.

6. (currently amended) A monocyclopentadienyl complex as claimed in claim 1 ~~any of claims 1 to 5~~, wherein Z is selected from the group consisting of $-C(R^{1B}R^{2B})-$, $Si(R^{3B}R^{4B})-$, $-CH_2-C(R^{3B}R^{4B})-$ and 1,2-phenylene.

7. (currently amended) A catalyst system for olefin polymerization comprising
 - A) at least one monocyclopentadienyl complex as claimed in claim 1 ~~claims 1 to 6~~,
 - B) optionally, an organic or inorganic support,
 - C) optionally, one or more activating compounds,
 - D) optionally, further catalysts suitable for olefin polymerization and
 - E) optionally, one or more metal compounds containing a metal of group 1, 2 or 13 of the Periodic Table.
8. (original) A prepolymerized catalyst system comprising a catalyst system as claimed in claim 7 and one or more linear C₂-C₁₀-1-alkenes polymerized onto it in a mass ratio of from 1: 0.1 to 1: 1 000 based on the catalyst system.
9. (currently amended) The use of a catalyst system as claimed in claim 7 ~~or 8~~ for the polymerization or copolymerization of olefins.
10. (currently amended) A process for preparing polyolefins by polymerization or copolymerization of olefins in the presence of a catalyst system as claimed in claim 7 ~~or 8~~.
11. (original) A process for preparing cyclopentadienyl systems of the formula (V):



where the variables have the following meanings:

$R^{1A}-R^{4A}$ are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part, NR^{6A}_2 , $N(SiR^{6A}_3)_2$, OR^{6A} , $OSiR^{6A}_3$, SiR^{6A}_3 , where the organic radicals $R^{1A}-R^{4A}$ may also be substituted by halogens and two vicinal radicals $R^{1A}-R^{4A}$ may also be joined to form a five- or six-membered ring, and/or two vicinal radicals $R^{1A}-R^{4A}$ are joined to form a heterocycle which contains at least one atom from the group consisting of N, P, O and S[[.]] ,

R^{6A} are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two geminal radicals R^{6A} may also be joined to form a five- or six-membered ring,

A is an unsubstituted, substituted or fused, heteroaromatic ring system,

$R^{1B}-R^{4B}$ are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part or SiR^{7B}_3 , where the organic radicals $R^{1B}-R^{4B}$ may also be substituted by halogens and two geminal vicinal radicals $R^{1B}-R^{4B}$ may also be joined to form a five- or six- membered ring and

R^{7B} are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl or alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two radicals R^{7B} may also be joined to form a five- or six-membered ring,


M^S a metal of group 1, 2 or 3 of the Periodic Table of the Elements,

X^S are each, independently of one another, fluorine, chlorine, bromine, iodine, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl, alkylaryl having 1-10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part, NR^1R^2 , OR^1 , SR^1 , SO_3R^1 , $OC(O)R^1$, CN , SCN , β -diketonate, CO , BF_4^- , PF_6^- or a bulky noncoordinating anion and


s 0, 1 or 2,

r 1 or 2, with the proviso that $s + r$ is the oxidation state of $M^S - 1$,

which comprises reacting $(A-CR^{1B}R^{2B})_r (M^S X^S)^+$ with a fulvene of the formula (VI)



12. (original) A process for preparing cyclopentadienyl systems of the formula (VIII):



where the variables have the following meanings:

$E^{6A}-E^{10A}$ are each carbon or not more than one E^{6A} to E^{10A} is phosphorus, where four adjacent $E^{6A}-E^{10A}$ form a conjugated diene system and the remaining $E^{6A}-E^{10A}$ additionally bears a hydrogen atom,

$R^{1A}-R^{4A}$ are each, independently of one another, hydrogen, C_1-C_{20} -alkyl, C_2-C_{20} -alkenyl, C_6-C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part, NR^{6A}_2 , $N(SiR^{6A}_3)_2$, OR^{6A} , $OSiR^{6A}_3$, SiR^{6A}_3 , where the organic radicals $R^{1A}-R^{4A}$ may also be substituted by halogens and two vicinal radicals $R^{1A}-R^{4A}$ may also be joined to form a five- or six-membered ring, and/or two vicinal radicals $R^{1A}-R^{4A}$ are joined to form a heterocycle which contains at least one atom from the group consisting of N, P, O and S,


R^{6A} are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two geminal radicals R^{6A} may also be joined to form a five- or six-membered ring,

A is an unsubstituted, substituted or fused, heteroaromatic ring system,

R^{1A} - R^{4A} are each, independently of one another, hydrogen C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl, alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part or SiR^{7B}_3 , where the organic radicals R^{1A} - R^{4A} may also be substituted by halogens and two geminal or vicinal radicals R^{1A} - R^{4A} may also be joined to form a five- or six-membered ring, and

R^{7B} are each, independently of one another, hydrogen, C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_6 - C_{20} -aryl or alkylaryl having from 1 to 10 carbon atoms in the alkyl part and 6-20 carbon atoms in the aryl part and two radicals R^{7B} may also be joined to form a five- or six-membered ring,

which comprises reacting $(A-CR^{1B}R^{2B})_r(M^S X^S)_s^+$ with a cyclopentadienyl system of the formula (IX)



where the variables are as defined above and

Q is a leaving group.